

Empirical Rules: If and only if the **boxplot** or **histogram** is **centered** then we can apply the **three** following empirical rules.

$68\% = \bar{x} \pm 1s$	68 % of data are within $1s$ of the mean (\bar{x})	
$95\% = \bar{x} \pm 2s$	95 % of data are within $2s$ of the mean (\bar{x})	
$99.7\% = \bar{x} \pm 3s$	99.7 % of data are within $3s$ of the mean (\bar{x})	

Example: Find all three empirical rules for Abe Stat final exam if the average was 72 and the standard deviation was 8, assuming that boxplot was centered.

$$68\% = 72 \pm 1(8) = 72 \pm 8$$

$$64 < \mathbf{68\%}$$
 of class got scores < 80

$$95\% = 72 \pm 2(8) = 72 \pm 16$$

$$56 < \mathbf{95\%}$$
 of class got scores < 88

$$99.7\% = 72 \pm 3(8) = 72 \pm 24$$

$$48 < \mathbf{99.7\%}$$
 of class got scores < 96

Extra Practice: Answer questions **C** on **page 3** of practice problem **part 1**

Z-score: is used to show the **relative position of a data point** with respect of the rest of data by **measuring how many standard deviation** that point is **away from the mean**. To apply the z-score the boxplot or histogram must be centered.

$$Z = \frac{x - \bar{x}}{s} \quad \text{or} \quad Z = \frac{x - \mu}{\sigma}$$

The possible range of Z-values.

If Z-value is **less than -2** or **more than 2** , it is called **unusual** that could be **unusually low** or **unusually high**.

If Z-value is **between -2 and 2** , then it is called **ordinary** or **common**.

----- -2 ----- 0 ----- 2 -----

Unusual Values:

Ordinary Values:

Unusual Values:

Example 1: Find the z-score of final exam for Tommy Yank in stat class at CSUS, if his score was 87, when the class average was 72 and the standard deviation was 8.

$$Z = \frac{x - \mu}{\sigma} = \frac{87 - 72}{8} = \frac{15}{8} = 1.875$$

Ordinary or **Unusual Value?**

So, he does relatively an ordinary performance relative to the rest of his class.

Example 2: Find the z-score of final exam for Marcy Tank in stat class at UC Davis, if his score was 82, when the class average was 71 and the standard deviation was 4.

$$Z = \frac{x - \mu}{\sigma} = \frac{82 - 71}{4} = \frac{11}{4} = 2.75$$

Ordinary or **Unusual Value?**

So, she does relatively better than the rest of her class and her z score is unusual.

Basic Probability

Probability of an event $E = P(E) = \frac{f}{n} = \frac{\text{The number of **desired** (success) **outcomes**}}{\text{The total number of **possible outcomes**}}$

$0 \leq P(E) \leq 1$

0	.25	.5	.75	1
Impossible	unlikely	even chance	likely	certain


If the **probability** of occurrence of an event such as event E is between $0 \leq P(E) < 5\%$ then its occurrence is called unusual.

Definition	<u>Examples</u>		
An experiment is an action, or trial, through which specific results (outcomes) are obtained.	Tossing a coin	Rolling a Die	Draw one card from deck of 52 cards
sample space = n All possible outcomes of an experiment are called	n = 2 sides (H,T) n = 2 outcomes	n = 6 sides (1,2,3,4,5,6) n = 6 outcomes	n = 52 cards n = 52 outcomes
Out of sample space how many is/are the desired outcome or outcomes ? That will be = f .	a tail f = 1 tail	an odd number (1,3,5) f = 3 odd numbers	an Ace f = 4 Aces
Probability is the measure of how likely an event to occur = $P(E) = f / n$	$P(T) = 1/2 = 50\%$	$P(\text{odd number}) = 3/6 = 50\%$	$P(\text{Ace}) = 4/52 = 1/13$

Three Types of Probability

- **Classical:** (equally probable outcomes). Like flipping a coin, rolling a die, drawing one card from a deck of cards. In this type of probabilities, we know the probability of getting for number 5 is always 1/6.
- **Empirical:** We need data like example A on page 2. So based on **available data**, the answer may be different each time. $P(E) = \frac{\text{Frequency of event } E}{\text{Total frequency}} = \frac{f}{n}$
- **Subjective:** Guess or intuition feelings (doctor feels patient has 80% chance of recovery).

Example A:



The sample space has **eight possible outcomes**, $\{O+, O-, A+, A-, B+, B-, AB+, AB-\}$

Example B: (an example of **empirical** probability): the outcomes may vary from sample to sample
 Frequency distribution of annual income for U.S. families

Income	Frequency (1000s)
Under \$10,000	5,216
\$10,000–\$14,999	4,507
\$15,000–\$24,999	10,040
\$25,000–\$34,999	9,828
\$35,000–\$49,999	12,841
\$50,000–\$74,999	14,204
\$75,000 & over	12,961
	69,597


Part 1: Find the probability that a randomly selected person from this group makes \$75,000 and over

- 1) Experiment: randomly selecting a person. 2) Sample space = $n = 69,597$
 3) His/her income is \$75,000 and over: $f = 12,961$ 4) Prob (\$75,000 and over) = $12,961 / 69,597 = 18.63\%$

Part 2: Find the probability that a randomly selected person from this group makes \$24,999 or less

- 1) Experiment: randomly selecting a person. 2) Sample space = $n = 69,597$
 3) His/her income is \$75,000 and over: $f = 19,763$ 4) Prob (\$24,999 or less) = $19,763 / 69,597 = 28.40\%$

Example C. : (an example of **empirical** probability)



What is the probability that you spin the dial on the left spinner, and you get yellow? $P(\text{yellow}) = 1/5$
 What is the probability that you spin the dial on the right spinner, and you get lose turn? $P(\text{Lose Turn}) = 1/12$

Example D (an example of **classical** probability)

In a deck of 52 cards there are 13 diamonds and 12 faces, and 4 aces. If one card is drawn randomly find the probability that

Solution:

- a) $P(\text{diamond}) = 13 / 52 = 25\%$ b) $P(\text{face}) = 12 / 52 = 23.08\%$ c) $P(\text{not face}) = 40 / 52 = 76.92\%$
 d) $P(\text{not diamond}) = 39 / 52 = 75\%$ e) $P(\text{diamond and face}) = 3 / 52 = 5.77\%$

Example E: (an example of **classical** probability)

If we roll 2 dice, then there are 36 possible outcomes meaning that the **sample space is 36** or $n = 36$

2	3	4	5	6	7
3	4	5	6	7	8
4	5	6	7	8	9
5	6	7	8	9	10
6	7	8	9	10	11
7	8	9	10	11	12

Solution:

- a) find the probability that the sum of rolling two dice is 10
Event or desired outcomes: a sum of 10 $\Rightarrow \{(4,6),(5,5),(6,4)\} \Rightarrow f = 3$
 Prob (a sum of 10) = $3/36 = 1/12 = 8.33\%$

- b) find the probability that the sum of rolling two dice is 7
Event or desired outcomes: a sum of 7 $\Rightarrow \{(1,6),(2,5),(3,4),(4,3),(5,2),(6,1)\} \Rightarrow f = 6$
 Prob (a sum of 10) = $6/36 = 1/6 = 16.67\%$


- c) find the probability that the sum of rolling two dice is not 7
Event or desired outcomes: a sum of not 7 $\Rightarrow \neq \{(1,6),(2,5),(3,4),(4,3),(5,2),(6,1)\} \Rightarrow f = 30$
 Prob (a sum of 10) = $30/36 = 5/6 = 83.33\%$

- d) find the probability that the sum of rolling is 10 or more
Event or desired outcomes: to get a sum 10 or more $\Rightarrow \{(4,6),(5,5),(5,6),(6,4),(6,5),(6,6)\} \Rightarrow f = 6$
 Prob (a sum of 10 or more) = $6/36 = 1/6 = 16.67\%$

- e) find the probability that their sum is 5
Event or desired outcomes: to get a sum of 5 $\Rightarrow \{(1,4),(2,3),(3,2),(4,1)\} \Rightarrow f = 4$
 Prob (a sum of 5) = $4/36 = 1/9 = 11.11\%$

Law of Large Numbers

- As an experiment is repeated over and over, the empirical probability of an event approaches the theoretical (actual) probability of the event.



Multiplication Rule (Keywords: and, both, all)

$$P(A \text{ and } B \text{ and } C \text{ and } \dots) = P(A)P(B)P(C)\dots$$

We use multiplication rule to find the probability that events A, B, C happen together or one after each other.

Hint:

When you make a selection out of a group by using multiplication rule be aware of **with** or **w/o** replacement effect.

In a deck of 52 cards there are 13 diamonds and 12 faces, and 4 aces.

If 2 cards are randomly drawn **w/o replacement**, what is the probability that both are diamonds?

$$P(\text{both diamond}) = \frac{13}{52} \cdot \frac{12}{51} = 5.88\%$$

If 2 cards are randomly drawn **with replacement**, what is the probability that both are diamonds?

$$P(\text{both diamond}) = \frac{13}{52} \cdot \frac{13}{52} = 6.25\%$$

There are 13 diamonds and 12 faces, and 4 aces in a deck of card.

If 4 cards are randomly drawn **w/o replacement** then,

a) What is the probability that all 4 are diamond and how likelihood is this?

$$\begin{array}{|c|} \hline 13 \text{ Diamo} \\ \hline 39 \text{ Others} \\ \hline 52 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 12 \text{ Diamo} \\ \hline 39 \text{ Others} \\ \hline 51 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 11 \text{ Diamo} \\ \hline 39 \text{ Others} \\ \hline 50 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 10 \text{ Diamo} \\ \hline 39 \text{ Others} \\ \hline 49 \\ \hline \end{array} \quad \frac{13}{52} \cdot \frac{12}{51} \cdot \frac{11}{50} \cdot \frac{10}{49} = .26\% \text{ very unlikely}$$

b) What is the probability that all 4 are aces and how likelihood is this?

$$\begin{array}{|c|} \hline 4 \text{ Aces} \\ \hline 48 \text{ Others} \\ \hline 52 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 3 \text{ Aces} \\ \hline 48 \text{ Others} \\ \hline 51 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 2 \text{ Aces} \\ \hline 48 \text{ Others} \\ \hline 50 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 1 \text{ Aces} \\ \hline 48 \text{ Others} \\ \hline 49 \\ \hline \end{array} \quad \frac{4}{52} \cdot \frac{3}{51} \cdot \frac{2}{50} \cdot \frac{1}{49} = .000369\% \text{ very much unlikely}$$

c) What is the probability that all 4 are faces and how likelihood is this?

$$\begin{array}{|c|} \hline 12 \text{ Faces} \\ \hline 40 \text{ Others} \\ \hline 52 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 11 \text{ Faces} \\ \hline 40 \text{ Others} \\ \hline 51 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 10 \text{ Faces} \\ \hline 40 \text{ Others} \\ \hline 50 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 9 \text{ Faces} \\ \hline 40 \text{ Others} \\ \hline 49 \\ \hline \end{array} \quad \frac{12}{52} \cdot \frac{11}{51} \cdot \frac{10}{50} \cdot \frac{9}{49} = .001828 = .1828\% \text{ much unlikely}$$

d) What is the probability that all 4 are non faces and how likelihood is this?

$$\begin{array}{|c|} \hline 12 \text{ faces} \\ \hline 40 \text{ non-faces} \\ \hline 52 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 12 \text{ faces} \\ \hline 39 \text{ non-faces} \\ \hline 51 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 12 \text{ faces} \\ \hline 38 \text{ non-faces} \\ \hline 50 \\ \hline \end{array} \Rightarrow \begin{array}{|c|} \hline 12 \text{ faces} \\ \hline 37 \text{ non-faces} \\ \hline 49 \\ \hline \end{array} \quad \frac{40}{52} \cdot \frac{39}{51} \cdot \frac{38}{50} \cdot \frac{37}{49} = .337575 = 33.76\% \text{ It is likely}$$

A. If we have a group of 4 men and 6 women, and we select two at random, without replacement, then

1. Find the probability that both are women. $P(\text{both W}) = P(\text{W and W}) = \frac{6}{10} \cdot \frac{5}{9} = \frac{30}{90} = 0.33$

2. Find the probability that one of each gender is selected. That means **one man one woman** or **one woman one man**

$$P(\text{MW}) = \frac{4}{10} \cdot \frac{6}{9} = \frac{24}{90} = \frac{8}{30} = 0.267 \quad \text{or} \quad P(\text{WM}) = \frac{6}{10} \cdot \frac{4}{9} = \frac{24}{90} = \frac{8}{30} = 0.267$$

Then you need to add these probabilities. $0.267 + 0.267 = 0.533 = 53.33\%$

B. In a bag there are 3 red, 4 blue and 5 green marbles. If we draw 3 marbles at random (**without replacement**) then,

Find the probability that all

1) All red $P(\text{RRR}) = \frac{3}{12} \cdot \frac{2}{11} \cdot \frac{1}{10} = \frac{1}{220} = 0.0004545$

2) Non red $\frac{9}{12} \cdot \frac{8}{11} \cdot \frac{7}{10} = \frac{21}{55} = 0.38181$

3) All blue $\frac{4}{12} \cdot \frac{3}{11} \cdot \frac{2}{10} = \frac{1}{55} = 0.018181$

4) None blue $\frac{8}{12} \cdot \frac{7}{11} \cdot \frac{6}{10} = \frac{14}{55} = 0.2545$

C. In a bag there are 3 red, 4 blue and 5 green marbles. If we draw 3 marbles at random (**with replacement**, then,

Find the probability that all

1) All red $P(\text{RRR}) = \frac{3}{12} \cdot \frac{3}{12} \cdot \frac{3}{12} = \frac{1}{64} = 0.0156$

2) Non red $\frac{9}{12} \cdot \frac{9}{12} \cdot \frac{9}{12} = \frac{27}{64} = 0.4219$

3) All blue $\frac{4}{12} \cdot \frac{4}{12} \cdot \frac{4}{12} = \frac{1}{27} = 0.037$

4) None blue $\frac{8}{12} \cdot \frac{8}{12} \cdot \frac{8}{12} = \frac{8}{27} = 0.2963$